

Geostatistics of Dependent and Asymptotically Independent Extremes

A. C. Davison, R. Huser and E. Thibaud

Received: date / Revised version: date

Abstract Spatial modeling of rare events has obvious applications in the environmental sciences and is crucial when assessing the effects of catastrophic events, such as heatwaves or widespread flooding, on food security and on the sustainability of societal infrastructure. Although classical geostatistics is largely based on Gaussian processes and distributions, these are not appropriate for extremes, for which max-stable and related processes provide more suitable models. This paper provides a brief overview of current work on the statistics of spatial extremes, with an emphasis on the consequences of the assumption of max-stability. Applications to winter minimum temperatures and daily rainfall are described.

Some key words: Asymptotic independence; Brown–Resnick process; Extreme-value statistics; Gaussian process; Generalised Pareto distribution; Max-stable process; Statistics of extremes

1 Introduction

1.1 Motivation

In recent years there has been a major upsurge of research activity in the statistics of extreme events for spatial settings. One reason for this is the realisation among stakeholders, such as climate scientists, environmental engineers and insurance companies, that in an evolving climate it may be

This paper is based on an invited lecture at GeoEnv2012. We thank the organisers, and particularly Jaime Gómez-Hernández, for their splendid hospitality, and reviewers for helpful comments on an earlier version of this paper. The work was supported by the Swiss National Science Foundation and by the ETH domain Competence Center Environment and Sustainability (<http://www.cces.ethz.ch/>).

Ecole Polytechnique Fédérale de Lausanne, EPFL-FSB-MATHAA-STAT, Station 8, 1015 Lausanne, Switzerland (<http://stat.epfl.ch>).

changes in the sizes and frequencies of rare events, rather than changes in the averages, that lead to the most devastating losses of life, damage to infrastructure and so forth. While it is difficult or even impossible to attribute particular events to the effect of climatic change, the types of events that have long been forecast to increase in frequency by the modeling community—such as heatwaves leading to crop failure and major brush fires, or heavy summer rainfall leading to widespread flooding—do indeed seem to be appearing more often than in the recorded past. This motivates attempts to model such events, in order to understand their likely future impacts, and to assess the related risks.

Classical geostatistics is a well-developed field surveyed in numerous textbooks (e.g., Cressie, 1993; Wackernagel, 2003; Banerjee *et al.*, 2004; Diggle and Ribeiro, 2007; Cressie and Wikle, 2011), with much software available and a wide range of user communities corresponding to its many applications. Its basis in Gaussian distributions makes it unsuitable for extremal modeling, however, because the Gaussian density function has an exceptionally light tail and therefore can badly underestimate probabilities associated to extreme events. Moreover the tails of the multivariate Gaussian distribution lead to independent extremes, for any underlying correlation that is less than unity, resulting in potentially disastrous underestimation of the probabilities of the simultaneous occurrence of two rare events—this is ‘the formula that killed Wall Street’ which, at least according to *Wired* magazine (Salmon, 2009), has played a key role in the ongoing international financial crisis by providing wildly incorrect assessment of economic risks.

1.2 Univariate extremes

Since Gaussian densities do not provide suitable models for extremes, it is natural to ask what distributions can arise as limits for maxima of independent random variables. An argument due to Fisher and Tippett (1928) and subsequently greatly extended (Leadbetter *et al.*, 1983; Galambos, 1987; Resnick, 2006; de Haan and Ferreira, 2006) shows that the suitably rescaled maxima of independent random variables, and of a wide variety of random processes, follow the generalized extreme-value distribution (GEV),

$$H(y) = \begin{cases} \exp \left[- \left\{ 1 + \xi(y - \mu)/\tau \right\}_+^{-1/\xi} \right], & \xi \neq 0, \\ \exp \left[- \exp \{ -(y - \mu)/\tau \} \right], & \xi = 0, \end{cases} \quad (1)$$

where $a_+ = \max(a, 0)$, $\mu \in \mathbb{R}$, $\tau > 0$ and $\xi \in \mathbb{R}$. The quantities μ and τ in equation (1) are respectively location and scale parameters, and the shape parameter ξ determines the weight of the upper tail of the density, with increasing ξ corresponding to higher probabilities of large events. Environmental data rarely display $|\xi| > 1$, and often estimates of ξ lie in the interval $(-1/2, 1/2)$, but data from finance and telecommunications applications may have heavier tails, with $\xi > 1$. The value of ξ has implications

for inference, since the r th moment of (1) exists only if $\xi < 1/r$, and the regularity conditions for standard likelihood inference hold only for $\xi > -1/2$. Thus moment- or likelihood-based inference may perform poorly, and alternative approaches, such as probability-weighted moments (Hosking *et al.*, 1985; Hosking and Wallis, 1987) or penalised likelihood (Martins and Steingard, 2000), have been promoted for use in simple settings. However in more complex applications such as those described below, their flexibility and generality have made likelihood and Bayesian approaches essential.

Expression (1) provides the natural class of models for maxima, from which corresponding models for other definitions of extremes, such as exceedances over high thresholds, may be derived. To be more precise, if extremes of a stream of independently identically distributed random variates are defined in terms of exceedances over a threshold u , then under the rescaling that yields (1) as the distribution of a maximum and a suitable time rescaling, exceedances of u arise as a Poisson process on the set $[0, 1] \times [u, \infty)$ with measure

$$\Lambda\{[t_1, t_2] \times [x, \infty)\} = (t_2 - t_1) \times \{1 + \xi(x - \mu)/\tau\}_+^{-1/\xi}, \quad x > u, \xi \neq 0, \quad (2)$$

and with the same parameters as in (1); when $\xi = 0$ the second factor on the right of (2) is replaced by $\exp\{-(x - \mu)/\tau\}$. This implies that the times of exceedances of u occur according to a Poisson process of constant rate $\lambda = \{1 + \xi(u - \mu)/\tau\}_+^{-1/\xi}$, and that conditional on there being an exceedance $X > u$ at time t , we have

$$\Pr(X > x + u \mid X > u) = \begin{cases} (1 + \xi x/\sigma)_+^{-1/\xi}, & \xi \neq 0, \\ \exp(-x/\sigma), & \xi = 0, \end{cases} \quad x > 0, \quad (3)$$

where $\sigma = \tau + \xi u$ is a scale parameter, and ξ takes the same value as in (1), so that $\xi = 0$ corresponds to an exponential distribution of exceedances. Expression (3) is the survivor function of the generalized Pareto distribution (GPD). This provides a semiparametric approximation to the distribution function underlying a random sample, whereby the empirical distribution function of the data is used below the threshold u , and the generalized Pareto distribution is fitted to those observations that exceed u . This involves the estimation of three parameters: the probability of exceeding u , and the GPD parameters σ and ξ .

Equation (1) satisfies the max-stability property: for any $m \in \mathbb{N}$ there exist real numbers $a_m > 0$ and b_m such that

$$H^m(b_m + a_m x) = H(x), \quad x \in \mathbb{R}. \quad (4)$$

This necessary condition for a limiting distribution for maxima is satisfied only by the GEV, giving it strong mathematical support as a suitable distribution for fitting to maxima of scalar random variables. A corresponding notion of threshold stability gives cogent arguments for fitting of the GPD to exceedances over high thresholds.

Lower extrema are treated simply by noting that $\min(X_1, \dots, X_n) = -\max(-X_1, \dots, -X_n)$: the results for maxima or for exceedances over a threshold are simply applied to a transformed version of the original variables, followed by a back-transformation.

If the observations are identically distributed but not independent, then under fairly mild conditions the renormalised maxima continue to follow distribution (1) with the same value of ξ , but threshold exceedances occur as a compound Poisson process, with clusters of mean size θ^{-1} , where $\theta \in (0, 1]$ is known as the extremal index. The limiting distribution of cluster maxima remains (3), and this is the same distribution as that of an arbitrary exceedance. For more details see Leadbetter *et al.* (1983), Hsing (1987), Hsing *et al.* (1988) and Leadbetter (1991).

1.3 Outline

Statistical models derived from equations (1), (2) and (3) have been widely used in applications for decades (Gumbel, 1958; Smith, 1989; Davison and Smith, 1990; Coles, 2001; Beirlant *et al.*, 2004), but modeling in the spatial and space-time domains is much more recent. It rests on extensions of equation (1) that satisfy an appropriate generalization of equation (4). In Section 2 we outline the construction of max-stable processes, and then sketch how inference for them is conducted. Other reviews, also involving comparisons with other approaches, include Davison *et al.* (2012) and Cooley *et al.* (2012a).

Max-stable processes $Z(x)$ defined on a spatial domain \mathcal{X} are asymptotically dependent: the conditional probability distribution of their values at two spatial locations, or sites, x_1 and x_2 , that differ by a lag vector h , satisfies

$$\chi_h = \lim_{z \rightarrow \infty} \Pr\{Z(x_1) > z \mid Z(x_2) > z\}, \quad x_1, x_2 \in \mathcal{X}, \quad (5)$$

where a strictly positive limit implies that a large event at x_2 leads to a non-zero probability of a similarly large event at x_1 . However in some applications it seems that $\chi_h = 0$, and then so-called asymptotic independence models may be preferred; Gaussian process models are asymptotically independent. In this case the rate at which the limit in (5) is approached is of critical importance. Apart from the Gaussian model, one class of such models consists of the so-called inverted max-stable models discussed in Section 3.1. In Section 3.2 we describe measures of dependence that are suitable either for asymptotic dependence or independence models, or to help to discriminate between them, and in Section 4 we discuss two applications of these different models for extremes in geosciences.

2 Max-stable processes

2.1 Basic ideas

Consider annual maxima of some phenomenon, such as temperature or daily rainfall, observed at a set of sites $\mathcal{D} = \{x_1, \dots, x_D\}$ within a spatial domain \mathcal{X} . It is natural to treat the corresponding observations $Z(x_1), \dots, Z(x_D)$ as stemming from a process $Z(x)$ defined for all $x \in \mathcal{X}$, and the natural model for each of these individually is given by equation (1), but with location, scale and shape parameters taken to be spatially-varying surfaces, giving $\mu(x)$, $\tau(x)$ and $\xi(x)$, for $x \in \mathcal{X}$. These surfaces may also depend on explanatory variables, such as distance from a coast or altitude. In applications they may be taken to be polynomials or use basis functions such as thin-plate splines, and their parameters may be estimated using likelihood methods. A closely related approach, typically used in Bayesian modelling using Markov chain Monte Carlo simulation, is to represent variation in extremal parameters through spatial Gaussian processes. For example, Cooley *et al.* (2007) fit rainfall exceedances using a GPD model (3) in which $\log \sigma$ follows such a process but the shape parameter has two values, and many similar models have appeared in the literature (e.g., Coles and Casson, 1998; Casson and Coles, 1999; Sang and Gelfand, 2009, 2010). Chavez *et al.* (2011) point out that inclusion of covariates in the usual parametrization of the GPD model leads to a lack of invariance to the choice of threshold that can be resolved using the GEV parametrization. One major difference between models such as that of Cooley *et al.* (2007) and those proposed below is that the former fit univariate extremal distributions to data at particular spatial locations, but treat the margins as independent, conditional on the covariates, so risk estimates for spatial quantities may be poor, though those at individual locations can be expected to improve because of borrowing of strength across the spatial domain. By contrast the models described here aim to capture joint spatial properties of extremes in addition to their marginal variation.

The exposition below is simplified if we assume that these marginal surfaces are known, in which case we can transform the original data so that $Z(x)$ has a unit Fréchet distribution, i.e., $\Pr\{Z(x) \leq z\} = \exp(-1/z)$, for $z > 0$ and each $x \in \mathcal{X}$.

The notion of max-stability extending (4) is that for each $m = 1, 2, \dots$, there exist continuous functions $a_m(x) > 0$ and $b_m(x)$ such that for suitable functions $z(x)$,

$$\Pr\{Z(x) \leq b_m(x) + a_m(x)z(x), x \in \mathcal{X}\}^m = \Pr\{Z(x) \leq z(x), x \in \mathcal{X}\}, \quad (6)$$

or, equivalently, $Z(x)$ and the maximum of m independent copies of $\{Z(x) - b_m(x)\}/a_m(x)$ have the same distribution. A continuity argument implies that this is true for any $m > 0$. The natural extension of the GEV is the class of max-stable processes, which, under mild conditions, comprise the only possible non-degenerate limits of linearly rescaled component-wise maxima of identically distributed processes (de Haan and Ferreira, 2006,

Chap. 9). They thus provide natural models for processes of maxima. Max-stable processes may be represented in the form (de Haan, 1984; Schlather, 2002)

$$Z(x) = \max_{i=1, \dots, \infty} W_i(x)/T_i, \quad x \in \mathcal{X}, \quad (7)$$

where $0 < T_1 < T_2 < \dots$ are the points of a unit rate Poisson process on \mathbb{R}_+ and the $W_i(x)$ are independent replicates of a positive continuous stochastic process $W(x)$ on \mathcal{X} that satisfies $\mathbb{E}\{W(x)\} \equiv 1$ and certain technical conditions. The representation (7) may be interpreted in terms of a “storm” process, where the $W_i(x)$ are storm shapes and the T_i^{-1} their intensities, and the largest storm observed at each site x is recorded.

Following Schlather (2002), it is possible to check that (7) satisfies (6) and has unit Fréchet marginal distributions, and that

$$\Pr\{Z(x) \leq z(x), x \in \mathcal{X}\} = \exp\left(-\mathbb{E}\left[\sup_{x \in \mathcal{X}} \left\{\frac{W(x)}{z(x)}\right\}\right]\right). \quad (8)$$

By setting $z(x_d) = z_d$ for $d = 1, \dots, D$ and $z(x) = +\infty$ elsewhere, we see that the joint cumulative distribution function of $Z(x_1), \dots, Z(x_D)$ equals $\exp\{-V(z_1, \dots, z_D)\}$, where the function

$$V(z_1, \dots, z_D) = \mathbb{E}\left[\max_{d \in \{1, \dots, D\}} \left\{\frac{W(x_d)}{z_d}\right\}\right] \quad (9)$$

is known as the exponent measure. This is easily shown to be homogeneous of order -1 , and must satisfy $V(\infty, \dots, z_d, \dots, \infty) = 1/z_d$ in order that each of the marginal distributions is unit Fréchet. To sketch one important implication of the homogeneity, let us write $z = z(x)$ and define the pseudo-polar coordinates $r = \|z\|$ and $w = z/\|z\|$, where $\|\cdot\|$ represents a norm. Then it can be shown that the exponent measure factorises as a product of two measures, one depending on the pseudo-radius r and the other on the pseudo-angle w . This factorization underpins extrapolation to high levels, since it implies that the angular component can be estimated from observed data, and then used to extrapolate beyond the observations, through the regularity of the radial component. See de Haan and Resnick (1977) and Beirlant *et al.* (2004, §8.2.3).

Expression (9) may be used to compute $V(z_1, \dots, z_D)$ for certain choices of $W(x)$, but typically it is only explicitly available for $D = 2$ (but see Genton *et al.*, 2011; Huser and Davison, 2013; Wadsworth and Tawn, 2013), so that full likelihood inference based on (8) seems unattainable in general; see §2.3.

For any pairs of sites $x_1, x_2 \in \mathcal{X}$, the strength of dependence between $Z(x_1)$ and $Z(x_2)$ may be summarized by the extremal coefficient $\theta(h) = V(1, 1) \in [1, 2]$, where $h = x_1 - x_2$ is the lag vector, with $\theta(h) = 1$ corresponding to perfect dependence and $\theta(h) = 2$ to independence. There is an equivalence with the limiting probability χ_h in (5), since $\chi_h = 2 - \theta(h)$, so $\chi_h = 1$ for perfectly dependent models, and $\chi_h = 0$ for independent and asymptotically independent cases. Thus one expects $\theta(h)$ to be monotonic




Fig. 1 Max-stable processes, with unit Fréchet margins, simulated from (a) the Smith model, for which $W(x)$ is a Gaussian density with covariance matrix whose diagonal and off-diagonal elements equal 0.1 and 0.05 respectively; (b) the Schlather model, $\varepsilon(x)$ having stable correlation function $\rho(h) = \exp(-2\|h\|)$.

increasing in $\|h\|$, analogous to the variogram of a spatial process. Analogues to $\theta(h)$ may be defined for more than two sites (Schlather and Tawn, 2003), and are useful for model-checking.

2.2 Models

Equation (7) and suitable choices of $W(x)$ can be used to construct max-stable models. The Gaussian extreme value process (Smith, 1990), often referred to as the Smith model, a realization of which is shown in the left panel of Fig. 1, is constructed by taking $W(x)$ to be a bivariate Gaussian density centred at a random point that is uniformly distributed in the spatial domain. This model seems too smooth to be useful in applications, and in this respect it is unfortunate that it was the first model to be fitted (Padoan *et al.*, 2010) and thus has in some sense become a standard.

Non-smooth processes are more suited for modeling natural phenomena. The Schlather (2002) model, a realization of which is shown in the right panel of Fig. 1, takes

$$W(x) = \sqrt{2\pi} \max\{0, \varepsilon(x)\}, \quad (10)$$

where $\varepsilon(x)$ is a stationary centred Gaussian process with unit variance and correlation function $\rho(h)$. The smoothness of the resulting process can be controlled by the choice of correlation function. However, the Schlather model is not ergodic in the sense that it cannot attain independence of $Z(x_1)$ and $Z(x_2)$ unless $\text{corr}\{\varepsilon(x_1), \varepsilon(x_2)\} = -1$; equivalently its extremal coefficient $\theta(h)$ is bounded below 2 for any h (Kabluchko and Schlather, 2010; Davison and Gholamrezaee, 2012). To circumvent this, Schlather (2002)

proposed to introduce a random set element that ensures that sites that are distant enough cannot be covered by the same random function $W_i(x)$ in (7) and thus yields exact independence between maxima at such sites.

Another possibility, which can be viewed as extending the Smith model (Huser and Davison, 2013), is the Brown–Resnick process (Brown and Resnick, 1977; Kabluchko *et al.*, 2009), sometimes called the geometric Gaussian process. This is constructed by using (7) with

$$W(x) = \exp\{\varepsilon(x) - \gamma(x)\}, \quad (11)$$

where $\varepsilon(x)$ is an intrinsically stationary Gaussian process with mean zero, semi-variogram $\gamma(h)$, and $\varepsilon(0) = 0$ almost surely. Brown–Resnick processes can be shown to be essentially the only limit of properly rescaled maxima of Gaussian processes (Kabluchko *et al.*, 2009) and have been found to yield good fits in applications (Davison *et al.*, 2012; Jeon and Smith, 2012). Their flexibility, and the recent development of efficient inference procedures (Engelke *et al.*, 2012; Wadsworth and Tawn, 2013), make them particularly attractive. Although Brown–Resnick processes are based on non-stationary Gaussian models, the construction in equation (7) ensures that their distributions are stationary and highly non-Gaussian. Furthermore, methods for simulating Brown–Resnick processes have been developed (Oesting *et al.*, 2012; Schlather, 2002; Ribatet, 2011), though they are more computationally intensive than for the Schlather or Smith models.

As for the univariate case with the GPD and the Poisson process of exceedances sketched in Section 1.2, the results for maxima can be generalized naturally to the modeling of threshold exceedances. Loosely speaking, it can be shown that under similar conditions, the dependence structure of very high spatial events, not necessarily maxima, converges to that of a max-stable process, so the models described above can also be applied for excesses of very large thresholds.

2.3 Inference

Owing to the complicated form of the density stemming from differentiation of (8) with (9), classical likelihood inference for the parameters of max-stable models is not possible in general, and recourse has been made to composite likelihoods (Varin *et al.*, 2011) based on lower-order marginal densities, such as bivariate margins of all pairs of maxima. Under mild conditions maximum composite likelihood estimators are strongly consistent and have asymptotic normal distributions, though they may be much more variable than ordinary maximum likelihood estimators (Huser and Davison, 2012; Davis and Yau, 2011). The composite likelihood information criterion, CLIC, the analogue of the AIC for composite likelihoods, allows model comparison (Varin and Vidoni, 2005; Davison and Gholamrezaee, 2012). More recently, inference methods based on the full likelihood have been proposed for a special class of max-stable processes that includes the Brown–Resnick

processes (Wadsworth and Tawn, 2013), and Engelke *et al.* (2012) proposed to fit Brown–Resnick processes based on the full likelihood of “extremal increments” of the process. From the Bayesian perspective, Ribatet *et al.* (2012) have developed a Monte Carlo Markov chain algorithm for fitting such models by sampling the pseudo composite posterior distribution using a modified acceptance rate. Reich and Shaby (2012) and Shaby and Reich (2012) describe the fitting of a Bayesian hierarchical model built from a finite approximation to a max-stable process, based on latent α -stable variates.

When individual events are recorded, more efficient inference is feasible. Since the max-stable models are suitable only above some predetermined high threshold, inference is usually made using a censored approach (Huser and Davison, 2012; Jeon and Smith, 2012; Huser and Davison, 2013; Thibaud and Davison, 2013). Furthermore, following Stephenson and Tawn (2005), Davison and Gholamrezaee (2012) and Wadsworth and Tawn (2013) show how to incorporate the occurrence times of extreme events, use of which both simplifies the likelihood and allows much more efficient inference in cases of moderate to low spatial dependence.

3 Asymptotic dependence and independence

3.1 Models for asymptotic independence

As mentioned in §1, stationary max-stable processes $Z(x)$ with unit Fréchet marginal distributions are asymptotically dependent in the sense that

$$\chi_h = \lim_{z \rightarrow \infty} \Pr\{Z(x_1) > z \mid Z(x_2) > z\} = 2 - \theta(h), \quad x_1, x_2 \in \mathcal{X}, \quad (12)$$

where $h = x_1 - x_2$ is the lag vector, and χ_h may be strictly positive. In practice it may be difficult to identify independence of extremes based on finite samples, since the data may display residual dependence for any finite threshold, however high. Asymptotic independence models, for which the limit in equation (12) equals zero, but which can also model the dependence present before the limit is reached, may therefore be preferred for modelling at finite thresholds. The Gaussian model is asymptotically independent for all correlations $\rho(h) \neq 1$, but Gaussian processes are too restrictive in the bulk of extremal applications, so broader classes of models are needed to allow flexible modeling.

Let $Z(x)$ denote a stationary process for spatial extremes with unit Fréchet margins. In order to smoothly link asymptotic dependence and independence in two dimensions, Ledford and Tawn (1996) proposed a model in which

$$\Pr\{Z(x_1) > z \mid Z(x_2) > z\} \sim \mathcal{L}(z)z^{1-1/\eta(h)}, \quad z \rightarrow \infty, \quad (13)$$

where the function $\mathcal{L}(z)$ is slowly varying at infinity, i.e., $\lim_{t \rightarrow \infty} \mathcal{L}(ta)/\mathcal{L}(t) = 1$, for any $a > 0$, and $\eta(h) \in (0, 1]$ is called the coefficient of tail dependence.

If $\eta(h) = 1$, then the limit in (13) depends on $\mathcal{L}(z)$ and is typically non-zero, cf. (12), whereas if $\eta(h) < 1$, then asymptotic independence arises: the dependence decreases with the size z of the rare event, at a rate determined by $\eta(h)$. In the case of a Gaussian process, $\eta(h) = \{1 + \rho(h)\}/2$. The variables are positively associated when $\eta(h) > 1/2$ and negatively associated when $\eta(h) < 1/2$. So-called near-independence corresponds to $\eta(h) = 1/2$, since in that case the conditional survivor function (13) corresponds to that of a Fréchet distribution. Further bivariate asymptotic independence models have been proposed, notably by Ledford and Tawn (1997) and Ramos and Ledford (2009, 2011).

It turns out that the lower tail of a multivariate max-stable distribution is asymptotically independent, and this motivated Wadsworth and Tawn (2012) to introduce the class of inverted max-stable processes, which are defined in terms of a max-stable process $Z(x)$ by

$$Z'(x) = -1/\log[1 - \exp\{-1/Z(x)\}], \quad x \in \mathcal{X}, \quad (14)$$

and which provide spatial models for asymptotic independence. For these processes, $\eta(h) = 1/\theta(h)$, where $\theta(h)$ is the extremal coefficient of $Z(x)$. With this construction, each max-stable model $Z(x)$ may be transformed to provide an asymptotically independent counterpart $Z'(x)$. Fig. 2 depicts conditional exceedance probabilities for max-stable, Gaussian and inverted max-stable processes, for various levels of dependence. The difference between these asymptotic dependence and asymptotic independence models is striking; for the former, the conditional probabilities are convex in z and converge to a positive value, whereas for the latter they are concave in z and tend to zero. The probabilities for the Gaussian and inverted max-stable processes tend very slowly to zero, and although both models are asymptotically independent they can be quite different at finite thresholds due to differences in the slowly varying functions in (13).

Wadsworth and Tawn (2012) proposed the use of hybrid models, i.e., mixtures of max-stable and inverted max-stable processes, that can assure asymptotic dependence at short distances and asymptotic independence at larger ones. As with max-stable processes, such models can be fitted using composite likelihood, and can be compared using the CLIC.

3.2 Measures of extremal dependence

Different measures of extremal dependence have been proposed. The extremal coefficient $\theta(h)$, introduced in Section 2, is suitable for asymptotically dependent processes, whose renormalised maxima converge to a non-trivial max-stable process. Empirical estimators of $\theta(h)$, based on the assumption of max-stability, have been suggested by Schlather and Tawn (2003) and Naveau *et al.* (2009), the latter based on the madogram, an analogue of the variogram for extremal processes. These estimators appear to yield satisfactory results within the max-stable framework, but cannot distinguish different degrees of asymptotic independence.




Fig. 2 Theoretical conditional exceedance probabilities $p(z) = \Pr\{Z(x_1) > z \mid Z(x_2) > z\}$ for (a) a max-stable process with $\theta(h) = 1, 1.2, 1.4, 1.6, 1.8, 2$ (from top to bottom); (b) a Gaussian process with correlation $\rho(h) = 0, 0.25, 0.75, 0.9, 0.95, 1$ (from bottom to top); (c) an inverted max-stable process with $\eta(h) = \{1 + \rho(h)\}/2$, and the same values of $\rho(h)$ as in (b) (also from bottom to top). The margins are uniform on $[0, 1]$.

The coefficient of tail dependence $\eta(h)$, introduced in section 3.1, is suited for asymptotically independent variables, and can be estimated by maximum likelihood (Ledford and Tawn, 1996) using $\min\{Z(x), Z(x+h)\}$. However, since the case $\eta(h) = 1$ corresponds to the entire class of max-stable processes, it provides no information about the strength of dependence of a given max-stable process. Moreover, discriminating between asymptotic dependence and asymptotic independence is difficult, because the dependence between variables may vanish very slowly as the level increases; see Fig. 2. Ledford and Tawn (1996) discuss testing for asymptotic independence.

Coles *et al.* (1999) suggested model-free diagnostics $\chi_h(u)$ and $\bar{\chi}_h(u)$ for distinguishing among these different types of tail dependence. If $Z(x_1)$ and $Z(x_2)$ are assumed to have uniform marginal distributions, then these coefficients can be expressed as

$$\chi_h(u) = 2 - \frac{\log C(u, u)}{\log u}, \quad \bar{\chi}_h(u) = \frac{2 \log(1 - u)}{\log \bar{C}(u, u)} - 1, \quad 0 \leq u \leq 1, \quad (15)$$

where C is the joint cumulative distribution function of $Z(x_1)$ and $Z(x_2)$, also called their copula (Nelsen, 2006), and

$$\bar{C}(u_1, u_2) = \Pr\{Z(x_1) > u_1, Z(x_2) > u_2\}$$

is the corresponding bivariate survivor function. The dependence measures in (15) are often estimated by their empirical rank-based counterparts, though the model-based estimators mentioned above will be more efficient, at least when the underlying model is reasonable. Note that $\lim_{u \rightarrow 1} \chi_h(u) = \chi_h$, defined in (5).

The coefficient $\chi_h(u)$, whose sign indicates whether the variables are positively or negatively associated for a given u (Nelsen, 2006), is equivalent




Fig. 3 The dependence measures $\chi_h(u)$ and $\bar{\chi}_h(u)$ for a max-stable process with $\theta(h) = 1.7$ (red), a Gaussian process with $\rho(h) = 0.7$ (blue) and an inverted max-stable process with $\eta(h) = \{1 + \rho(h)\}/2 = 0.85$ (green).

to the conditional probability $\Pr\{Z(x_1) > u \mid Z(x_2) > u\}$ when $u \rightarrow 1$. In case of asymptotic independence, $\lim_{u \rightarrow 1} \chi_h(u) = 0$, and $\chi_h(u) \equiv 0$ for exactly independent variables. If the variables are max-stable with extremal coefficient $\theta(h)$, then $\chi_h(u) \equiv 2 - \theta(h) \geq 0$.

The coefficient $\bar{\chi}_h(u)$ provides a measure of dependence within the class of asymptotically independent processes. One can show that under (13), $\bar{\chi}_h(u) \rightarrow 2\eta(h) - 1$, as $u \rightarrow 1$. Furthermore, for any inverted max-stable process, $\bar{\chi}_h(u) \equiv 2\eta(h) - 1$.

Fig. 3 displays $\chi_h(u)$ and $\bar{\chi}_h(u)$ for max-stable, Gaussian and inverted max-stable processes. For max-stable processes, $\chi_h(u)$ is constant and $\bar{\chi}_h(u)$ increases to unity; for Gaussian processes, $\chi_h(u)$ decreases to zero and $\bar{\chi}_h(u)$ increases to $2\eta(h) - 1$; and for inverted max-stable processes, $\chi_h(u)$ decreases to zero and $\bar{\chi}_h(u)$ takes the constant value $2\eta(h) - 1$. Hence, in this sense, the Gaussian model has even lighter tails than the left tail of a max-stable process.

Similar dependence measures for extremes of time series are described by Ledford and Tawn (2003) and Davis and Mikosch (2010).

4 Applications

The ideas outlined above have been applied to a variety of extremal problems, often with further explanatory variables used to account for variation in the surfaces $\mu(x)$, $\tau(x)$ and $\xi(x)$, though $\xi(x)$ is often found to be constant. Buishand *et al.* (2008) take a semiparametric approach to max-stable modeling for rainfall data in a homogeneous region of the Netherlands. Padoan *et al.* (2010) fit the Smith model to annual maximum daily rain-

fall at 46 sites in the Appalachian mountains, and overall find that their best max-stable model fits reasonably well. Blanchet and Davison (2011) use similar models, but with regional effects and a third dimension, namely altitude, in a study of maximum snow depth in the Swiss Alps. Davison *et al.* (2012) compare a variety of models for extreme rainfall, including some based on copulas, Bayesian hierarchical distributions and max-stable models, and conclude that the latter fit best, but that the Smith model is clearly beaten by rougher processes like those in the right-hand panel of Fig. 1. Davison and Gholamrezaee (2012) apply the Schlather model with a random set to data on extreme temperatures, a relatively large-scale phenomenon compared to rainfall, and Huser and Davison (2012) have extended this to space-time modeling of extreme rainfall. Max-stable processes appear to be well suited for modeling rainfall and temperature processes at various scales.

The use of inverted max-stable models is more recent. Wadsworth and Tawn (2012) show that max-stable processes are not well adapted for modeling wave height data in North Sea and that inverted models are preferred. Thibaud and Davison (2013) compare the fits of Gaussian, max-stable and inverted max-stable models for extreme spatial rainfall in a small mountain catchment, and conclude that the last fits their data best.

Below we discuss a small spatial application using data on air temperature and precipitation, and find that asymptotic dependence models may be preferred for the first dataset but that asymptotic independence models seem to fit better for the second. Gaussian models perform less well in both cases. We consider winter daily temperature minima and summer daily cumulative rainfall, recorded at eight monitoring sites with similar altitudes and located in the so-called plateau region of Switzerland; see Fig. 4. The data were available from 1981 to 2012, giving a total of about 2900 observations per site. For simplicity, we treat these daily data as independent over time, although this is false at least for the temperature data. A more complex spatio-temporal study was performed by Huser and Davison (2012), where further details about part of the rainfall data may be found. They fitted several space-time max-stable models to hourly rainfall, of which the Schlather model with random set seemed to be well-suited, but did not compare them to asymptotic independence models.

We first transformed the temperature data by multiplication by -1 , and then fitted the generalized Pareto distribution (3) to model events above the 98% quantile, u_{98} , of the time series at each site separately, and used this fitted model to transform the data to have the unit Fréchet distribution. Since Bortot *et al.* (2000) and Coles and Pauli (2002) have shown that the choice between asymptotic dependence and asymptotic independence models can influence extrapolation and association of extremes far more than the particular model used in each of these model classes, we fitted only a limited selection of spatial correlations for this illustrative analysis. For each dataset, we used a censored pairwise threshold-based approach (Huser and Davison, 2012) to fit three models to the exceedances over u_{98} :




Fig. 4 Topographic map of monitoring sites (with their altitudes) from *MeteoSwiss*, where temperature and precipitation data were recorded. The most distant sites are 107km apart, while the closest are 16km apart.

(a) the max-stable Brown–Resnick model in equation (11), with variogram $2\gamma(h) = \sigma^2 I(h > 0) + (\|h\|/\lambda)^\alpha$, where $\sigma > 0$ is a nugget effect, $I(\cdot)$ denotes the indicator function, $\lambda > 0$ is a range parameter and $\alpha \in (0, 2]$ is a smoothness parameter; (b) the corresponding inverted max-stable model; and (c) the multivariate Gaussian distribution, transformed to unit Fréchet margins, with correlation function $\rho(h) = \exp\{-\gamma(h)\}$. Larger values of α give smoother processes in each case. For the temperature data, α was always estimated very close to its upper boundary, so we set $\alpha = 2$ and estimated only the remaining parameters. Table 1 reports the estimated parameters and confidence intervals, and the corresponding values of the log pairwise likelihood and CLIC*, for each model and dataset. There is a strong difference in the scales and in the smoothness of the processes. The temperature data have $\hat{\lambda} = 210\text{km}$ or more, and $\alpha = 2$, corresponding to a relatively smooth process with large-scale dependence, though local variation is accommodated by the nugget σ^2 , which is significantly larger than zero. The rainfall data have $\hat{\lambda} > 3\text{km}$ and $0 < \hat{\alpha} < 1$, corresponding to a much rougher process, now with a nugget whose confidence interval almost includes zero. There is a clear trade-off between including a nugget to allow local variation in a max-stable process, and of using an asymptotically independent process, in which local variation can be stronger; the availability of data at nearby locations might allow better discrimination between these. In each case the value of $\hat{\lambda}$ is smallest for the max-stable process and seems unrealistically large for the Gaussian process. The rather wide confidence intervals for λ for the temperature data probably stem from the difficulty of estimating continental-scale events from data over a limited region of a small country.

Table 1 Estimated dependence parameters $\hat{\sigma}^2$, $\hat{\lambda}$ and $\hat{\alpha}$ (with 95% confidence intervals based on block bootstrap using seasonal blocks) from the models (a), (b) and (c), along with the value of the log pairwise likelihood, $\hat{\ell}$, evaluated at the parameter estimates and scaled in such a way that its value is comparable to the log likelihood under independence, as suggested by Davison and Gholamrezaee (2012). The values of the CLIC*, a scaled version of the CLIC, are also reported.

Dataset	Model	$\sigma^2 \times 10^2$	λ [km] $\times 10^{-1}$	α	$\hat{\ell}$	CLIC*
Temp.	(a) MS	58 (50, 71)	21 (15, 579)	2 (-)	-4289.6	8599.2 [†]
	(b) IMS	4 (3, 6)	77 (58, 668)	2 (-)	-4295.6	8604.5
	(c) Gauss	4 (3, 6)	80 (60, 3062)	2 (-)	-4291.8	8599.9
Rainfall	(a) MS	11 (0, 272)	0.3 (0.1, 3.7)	0.55 (0.36, 1.56)	-4836.9	9711.9
	(b) IMS	6 (0, 28)	11 (7, 23)	0.92 (0.52, 1.95)	-4829.4	9673.0 [‡]
	(c) Gauss	4 (0, 21)	31 (16, 111)	0.65 (0.38, 1.76)	-4829.9	9689.9

MS: max-stable model; IMS: inverted max-stable model; Gauss: Gaussian model.

[†] Our preferred model for the temperature data.

[‡] Our preferred model for the rainfall data.




Fig. 5 Extremal coefficient $\theta(h)$ (left) and coefficient of tail dependence $\eta(h)$ (right), corresponding to the best fitted models for the temperatures and rainfall data, respectively. According to Table 1, these models are respectively max-stable and inverted max-stable. Circles are binned empirical estimates with 95% confidence intervals in grey, while black solid lines are model-based estimates. The black points represent theoretical values of $\theta(0)$ and $\eta(0)$.

Fig. 5 displays binned empirical estimates of the extremal coefficient $\theta(h)$ for the temperature data, for which the best model is max-stable, and of the coefficient of tail dependence $\eta(h)$ for the rainfall data, which seem asymptotically independent. Comparing these estimates to their fitted counterparts, it seems that the models quite flexibly capture spatial dependence. The graphs confirm that the natural processes considered here have

decreasing extremal dependence with increasing distance, although dependence remains strong at long distances for temperatures, perhaps explaining the difficulty in estimating the range parameter. The huge uncertainty in these plots would be reduced by taking more sites. Moreover, given that the monitoring sites are at least 16km apart, small local variation represented by the nugget is difficult to estimate.

While the best model for the temperature data, according to the CLIC and the value of the log pairwise likelihood, is max-stable, the rainfall data appear to be asymptotically independent. Since max-stable models provide the only possible limiting dependence structure, the fit of these models should be better for higher thresholds, but as finite thresholds must be considered in practice, asymptotic independence models might provide better fits to the available data. If interest resides in extremely high joint return levels, it could be misleading to base inferences on asymptotic independence models. To assess this, Fig. 6 shows the empirical values for $\chi_h(u)$ and $\bar{\chi}_h(u)$ for the sites NEU and PAY, and their model-based counterparts. Although the fits at finite thresholds are similar for the different models, they differ at very extreme thresholds, and the probabilities of very extreme events might be strongly underestimated if an asymptotic independence model is used. The right panels of Fig. 6 display joint return levels for some spatial functionals of the daily temperature and rainfall data recorded at NEU and PAY. At observed timescales, the curves for the different models give more or less the same predictions, but at the 1000 year-return period, a discrepancy of around 7.6mm appears for the rainfall data, underlining the ability of the max-stable model to capture dependence at high levels. This effect would probably increase for longer return periods and if more sites were to be considered simultaneously. In summary, if there were any guarantees that the data we consider are indeed asymptotically independent, the Gaussian or inverted max-stable models would potentially be suitable, but since it is very difficult, if not impossible, to have insight so far into the tail, it is safer to base risk assessments on max-stable models, which provide upper bounds for joint probabilities of extreme events.

5 Discussion

This paper sketches some of the statistical ideas useful for modeling spatial extremes. Classical geostatistical models based on the Gaussian distribution may provide poor models for maxima or threshold exceedances. Since the multivariate normal distribution belongs to the class of asymptotically independent processes, it may underestimate the probabilities of rare events and thus provide poor risk estimates for spatial functionals. This difficulty also arises for inverted max-stable processes, presented in Section 3.1, even though these appear to be more flexible than the Gaussian distribution. Max-stable processes, which assume a degree of dependence that does not change with the severity of the event, are natural models to capture extremal




Fig. 6 Extremal diagnostics computed for the temperature (top row) and rainfall (bottom row) data for the pair of monitoring sites NEU and PAY. *Left–Middle columns*: Empirical estimates (circles) of $\chi_h(u)$ (left) and $\bar{\chi}_h(u)$ (middle) with their model-based counterparts: The solid lines show the fitted curves for (a) the max-stable model (red), (b) the inverted max-stable model (green) and (c) the Gaussian model (blue). The grey vertical lines are 95% confidence intervals from a block bootstrap using seasonal blocks, and the vertical dashed line is the threshold used when fitting the dependence models. *Right column*: Empirical and model-based return levels for the spatial average (over both sites) of daily temperature minima (top) and cumulative rainfall (bottom), on the original scale.

dependence. At finite thresholds these asymptotic dependence models may yield worse fits than flexible asymptotic independence models, but unless the data are truly asymptotically independent, which is difficult to assess in practice, max-stable processes provide a conservative basis for the estimation of probabilities of very rare events.

In order to discriminate between asymptotic independence and max-stability, different measures of extremal dependence have been introduced, including the diagnostics $\chi_h(u)$ and $\bar{\chi}_h(u)$ of Coles *et al.* (1999). Unlike the extremal coefficient and the coefficient of tail dependence, these two measures are nonparametric. They can be used to check whether the assumption of max-stability is reasonable, but must be interpreted with care at high thresholds.

Our data analysis suggests that daily temperature minima, which represent a large-scale phenomenon, tend to favor max-stable models, whereas daily rainfall totals show weaker spatial dependence and seem closer to asymptotic independence. Although spatial return levels at observed time-scales differ little, a discrepancy was visible for the rainfall data at the 1000 year-return period.

For space-time data, hybrid models (Wadsworth and Tawn, 2012) can capture different asymptotic properties at different distances and time lags, but they may be quite difficult to fit in practice. Coles and Pauli (2002) suggest an approach to merging the uncertainty arising from the dichotomy between the two types of asymptotic behaviour, but it is unclear how it might be applied in the spatial context.

An alternative approach is the use of conditional models for extremes, in which the behaviour of several variables is considered, conditioned on the event that one of them is extreme. Following its introduction by Heffernan and Tawn (2004), this approach has been developed from the theoretical viewpoint by Heffernan and Resnick (2007), Das and Resnick (2011) and Fougères and Soulier (2012), among others, and has been applied to spatial flood estimation by Keef *et al.* (2009, 2013). Related work is described by Cooley *et al.* (2012b).

Extrapolation to estimate probabilities of events never yet seen rests entirely on assumptions about the behaviour of distribution tails, but these can only be verified with respect to events that have already occurred. Thus the likely validity of the model underlying the extrapolation needs exceptionally careful consideration. This is compounded in the spatial setting, in which extrapolation from a finite and perhaps unreliable dataset to an infinite-dimensional space is required. This task at first appears impossible, but, if used with attention and supplemented with subject-matter knowledge, the ideas sketched in this paper provide initial steps towards a quantitative understanding of spatial rare events.

References

- Banerjee, S., Carlin, B. and Gelfand, A. (2004) *Hierarchical Modeling and Analysis for Spatial Data*. New York: Chapman and Hall/CRC.
- Beirlant, J., Goegebeur, Y., Teugels, J. and Segers, J. (2004) *Statistics of Extremes: Theory and Applications*. New York: Wiley.
- Blanchet, J. and Davison, A. C. (2011) Spatial modelling of extreme snow depth. *Annals of Applied Statistics* **5**, 1699–1725.
- Bortot, P., Coles, S. G. and Tawn, J. A. (2000) The multivariate Gaussian tail model: An application to oceanographic data. *Applied Statistics* **49**, 31–49.
- Brown, B. M. and Resnick, S. I. (1977) Extreme values of independent stochastic processes. *Journal of Applied Probability* **14**, 732–739.

- Buishand, T. A., de Haan, L. and Zhou, C. (2008) On spatial extremes: With application to a rainfall problem. *Annals of Applied Statistics* **2**, 624–642.
- Casson, E. and Coles, S. (1999) Spatial regression models for extremes. *Extremes* **1**, 449–468.
- Chavez, V., Davison, A. C. and Frossard, L. (2011) Discussion of “Threshold modelling of spatially-dependent non-stationary extremes with application to hurricane-induced wave heights” by P. J. Northrop and P. Jonathan. *Environmetrics* **22**, 810–812.
- Coles, S. (2001) *An Introduction to Statistical Modeling of Extreme Values*. London: Springer-Verlag.
- Coles, S., Heffernan, J. and Tawn, J. (1999) Dependence measures for extreme value analyses. *Extremes* **2**, 339–365.
- Coles, S. G. and Casson, E. (1998) Extreme value modelling of hurricane wind speeds. *Structural Safety* **20**, 283–296.
- Coles, S. G. and Pauli, F. (2002) Models and inference for uncertainty in extremal dependence. *Biometrika* **89**, 183–196.
- Cooley, D., Cisewski, J., Erhardt, R. J., Jeon, S., Mannshardt, E., Omolo, B. O. and Sun, Y. (2012a) A survey of spatial extremes: Measuring spatial dependence and modeling spatial effects. *REVSTAT—Statistical Journal* **10**, 135–165.
- Cooley, D., Davis, R. A. and Naveau, P. (2012b) Approximating the conditional density given large observed values via a multivariate extremes framework with application to environmental data. *Annals of Applied Statistics* **6**, 1406–1429.
- Cooley, D., Nychka, D. and Naveau, P. (2007) Bayesian spatial modeling of extreme precipitation return levels. *Journal of the American Statistical Association* **102**, 824–840.
- Cressie, N. A. C. (1993) *Statistics for Spatial Data*. New York: Wiley.
- Cressie, N. A. C. and Wikle, C. K. (2011) *Statistics for Spatio-Temporal Data*. New York: Wiley.
- Das, B. and Resnick, S. I. (2011) Conditioning on an extreme component: Model consistency with regular variation on cones. *Bernoulli* **17**, 226–252.
- Davis, R. A. and Mikosch, T. (2010) The extremogram: A correlogram for extreme events. *Bernoulli* **15**, 977–1009.
- Davis, R. A. and Yau, C. Y. (2011) Comments on pairwise likelihood in time series models. *Statistica Sinica* **21**, 255–277.
- Davison, A. C. and Gholamrezaee, M. M. (2012) Geostatistics of extremes. *Proceedings of the Royal Society of London, series A* **468**, 581–608.
- Davison, A. C., Padoan, S. A. and Ribatet, M. (2012) Statistical modelling of spatial extremes (with Discussion). *Statistical Science* **27**, 161–186.
- Davison, A. C. and Smith, R. L. (1990) Models for exceedances over high thresholds (with Discussion). *Journal of the Royal Statistical Society, series B* **52**, 393–442.
- de Haan, L. and Ferreira, A. (2006) *Extreme Value Theory: An Introduction*. New York: Springer.

- de Haan, L. and Resnick, S. I. (1977) Limit theory for multivariate sample extremes. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete* **40**, 317–337.
- Diggle, P. J. and Ribeiro, P. J. (2007) *Model-based Geostatistics*. New York: Springer.
- Engelke, S., Malinowski, A., Kabluchko, Z. and Schlather, M. (2012) Estimation of Hüsler–Reiss distributions and Brown–Resnick processes. arXiv:1207.6886v1 [stat.ME].
- Fisher, R. A. and Tippett, L. H. C. (1928) Limiting forms of the frequency distribution of the largest or smallest member of a sample. *Mathematical Proceedings of the Cambridge Philosophical Society* **24**, 180–190.
- Fougères, A.-L. and Soulier, P. (2012) Estimation of conditional laws given an extremal component. *Extremes* **15**, 1–34.
- Galambos, J. (1987) *The Asymptotic Theory of Extreme Order Statistics*. Second edition. Melbourne, FL: Krieger.
- Genton, M. G., Ma, Y. and Sang, H. (2011) On the likelihood function of Gaussian max-stable processes. *Biometrika* **98**, 481–488.
- Gumbel, E. J. (1958) *Statistics of Extremes*. New York: Columbia University Press.
- de Haan, L. (1984) A spectral representation for max-stable processes. *Annals of Probability* **12**, 1194–1204.
- Heffernan, J. E. and Resnick, S. I. (2007) Limit laws for random vectors with an extreme component. *Annals of Applied Probability* **17**, 537–571.
- Heffernan, J. E. and Tawn, J. A. (2004) A conditional approach for multivariate extreme values (with Discussion). *Journal of the Royal Statistical Society, series B* **66**, 497–546.
- Hosking, J. R. M. and Wallis, J. R. (1987) Parameter and quantile estimation for the generalized Pareto distribution. *Technometrics* **29**, 339–349.
- Hosking, J. R. M., Wallis, J. R. and Wood, E. F. (1985) Estimation of the generalized extreme-value distribution by the method of probability-weighted moments. *Technometrics* **27**, 251–261.
- Hsing, T. (1987) On the characterization of certain point processes. *Stochastic Processes and their Applications* **26**, 297–316.
- Hsing, T., Hüsler, J. and Leadbetter, M. R. (1988) On the exceedance point process for a stationary sequence. *Probability Theory and Related Fields* **78**, 97–112.
- Huser, R. and Davison, A. C. (2012) Space-time modelling of extreme events. arXiv:1201.3245v1 [stat.ME].
- Huser, R. and Davison, A. C. (2013) Composite likelihood estimation for the Brown–Resnick process. *Biometrika* **100**, to appear.
- Jeon, S. and Smith, R. L. (2012) Dependence structure of spatial extremes using threshold approach. arXiv:1209.6344v1 [stat.ME].
- Kabluchko, Z. and Schlather, M. (2010) Ergodic properties of max-infinitely divisible processes. *Stochastic Processes and their Applications* **120**, 281–295.

- Kabluchko, Z., Schlather, M. and De Haan, L. (2009) Stationary max-stable fields associated to negative definite functions. *The Annals of Probability* **37**, 2042–2065.
- Keef, C., Tawn, J. A. and Lamb, R. (2013) Estimating the probability of widespread floods. *Environmetrics* **24**, 13–21.
- Keef, C., Tawn, J. A. and Svensson, C. (2009) Spatial risk assessment for extreme river flows. *Applied Statistics* **58**, 601–618.
- Leadbetter, M. R. (1991) On a basis for ‘peaks over threshold’ modelling. *Statistics & Probability Letters* **12**, 357–362.
- Leadbetter, M. R., Lindgren, G. and Rootzén, H. (1983) *Extremes and Related Properties of Random Sequences and Processes*. New York: Springer-Verlag.
- Ledford, A. W. and Tawn, J. A. (1996) Statistics for near independence in multivariate extreme values. *Biometrika* **83**, 169–187.
- Ledford, A. W. and Tawn, J. A. (1997) Modelling dependence within joint tail regions. *Journal of the Royal Statistical Society, series B* **59**, 475–499.
- Ledford, A. W. and Tawn, J. A. (2003) Diagnostics for dependence within time series extremes. *Journal of the Royal Statistical Society, series B* **65**, 521–543.
- Martins, E. and Stedinger, J. (2000) Generalized maximum-likelihood generalized extreme-value quantile estimators for hydrologic data. *Water Resources Research* **36**, 737–744.
- Naveau, P., Guillou, A., Cooley, D. and Diebolt, J. (2009) Modelling pairwise dependence of maxima in space. *Biometrika* **96**, 1–17.
- Nelsen, R. B. (2006) *An Introduction to Copulas*. Second edition. New York: Springer.
- Oesting, M., Kabluchko, Z. and Schlather, M. (2012) Simulation of Brown–Resnick processes. *Extremes* **15**, 89–107.
- Padoan, S. A., Ribatet, M. and Sisson, S. A. (2010) Likelihood-based inference for max-stable processes. *Journal of the American Statistical Association* **105**, 263–277.
- Ramos, A. and Ledford, A. W. (2009) A new class of models for bivariate joint tails. *Journal of the Royal Statistical Society, series B* **71**, 219–241.
- Ramos, A. and Ledford, A. W. (2011) An alternative point process framework for modeling multivariate extreme values. *Communications in Statistics — Theory and Methods* **40**, 2205–2224.
- Reich, B. J. and Shaby, B. A. (2012) A hierarchical max-stable spatial model for extreme precipitation. *Annals of Applied Statistics* **6**, 1430–1451.
- Resnick, S. I. (2006) *Heavy-Tail Phenomena: Probabilistic and Statistical Modeling*. New York: Springer.
- Ribatet, M. (2011) *SpatialExtremes: Modelling Spatial Extremes*. R package version 1.8-5.
- Ribatet, M., Cooley, D. and Davison, A. C. (2012) Bayesian inference from composite likelihoods, with an application to spatial extremes. *Statistica Sinica* **22**, 813–845.

- Salmon, F. (2009) Recipe for disaster: The formula that killed Wall Street. http://www.wired.com/techbiz/it/magazine/17-03/wp_quant?currentPage=al, accessed on 6 March 2013.
- Sang, H. and Gelfand, A. E. (2009) Hierarchical modeling for extreme values observed over space and time. *Environmental and Ecological Statistics* **16**, 407–426.
- Sang, H. and Gelfand, A. E. (2010) Continuous spatial process models for spatial extreme values. *Journal of Agricultural, Biological and Environmental Statistics* **15**, 49–65.
- Schlather, M. (2002) Models for stationary max-stable random fields. *Extremes* **5**, 33–44.
- Schlather, M. and Tawn, J. A. (2003) A dependence measure for multivariate and spatial extreme values: Properties and inference. *Biometrika* **90**, 139–156.
- Shaby, B. A. and Reich, B. J. (2012) Bayesian spatial extreme value analysis to assess the changing risk of concurrent high temperatures across large portions of European cropland. *Environmetrics* **23**, 638–648.
- Smith, R. L. (1989) Extreme value analysis of environmental time series: An example based on ozone data (with Discussion). *Statistical Science* **4**, 367–393.
- Smith, R. L. (1990) Max-stable processes and spatial extremes. <http://www.stat.unc.edu/postscript/rs/spatex.pdf>.
- Stephenson, A. and Tawn, J. A. (2005) Exploiting occurrence times in likelihood inference for componentwise maxima. *Biometrika* **92**, 213–227.
- Thibaud, E. and Davison, A. C. (2013) Threshold modeling of extreme spatial rainfall. Submitted.
- Varin, C., Reid, N. and Firth, D. (2011) An overview of composite likelihood methods. *Statistica Sinica* **21**, 5–42.
- Varin, C. and Vidoni, P. (2005) A note on composite likelihood inference and model selection. *Biometrika* **92**, 519–528.
- Wackernagel, H. (2003) *Multivariate Geostatistics: An Introduction with Applications*. Third edition. New York: Springer.
- Wadsworth, J. L. and Tawn, J. A. (2012) Dependence modelling for spatial extremes. *Biometrika* **99**, 253–272.
- Wadsworth, J. L. and Tawn, J. A. (2013) Efficient inference for spatial extreme value processes associated to log-Gaussian random functions. Submitted to *Biometrika*.