

Discussion of “Visualizing spatiotemporal models with virtual reality: from fully immersive environments to applications in stereoscopic view” by Stefano Castruccio, Marc G. Genton and Ying Sun

Raphaël Huser¹, Miguel de Carvalho² and Luigi Lombardo¹

September 3, 2018

We congratulate the authors on this timely paper, which explores modern visualization tools for communicating statistical results and uncertainty more efficiently for spatiotemporal models defined on complex domains.

Although the authors chose to focus on classical space-time applications, the benefits of advanced visualization tools reach far beyond the standard Gaussian geostatistical setting. We highlight two different examples below of risk assessment applications.

Statistics of extremes ([Davison *et al.*, 2012](#); [Davison and Huser, 2015](#); [Davison *et al.*, 2018](#)) focuses on estimating low-probability tail events, and has obvious applications in Earth science and finance, where the notion of risk is often related to individual or simultaneous extreme events. [Genton *et al.* \(2015\)](#) used “visuanimations” to dynamically explore the dependence characteristics of a (highly non-Gaussian) spatiotemporal model for precipitation extremes proposed by [Huser and Davison \(2014\)](#). In this context, interactive visualization tools could be very useful, especially when adding an extra “tail” dimension, allowing one to dynamically explore the estimated joint tail behavior and efficiently visualize extreme spatiotemporal return levels for increasing return periods.

The most important object to estimate and visualize from point pattern data ([Cressie, 1993](#)), such as wild fire or landslide data, is their *intensity function*. [Lombardo *et al.* \(2018\)](#)

¹King Abdullah University of Science and Technology (KAUST), Computer, Electrical and Mathematical Sciences and Engineering (CEMSE) Division, Thuwal 23955-6900, Saudi Arabia. E-mail: raphael.huser@kaust.edu.sa

²University of Edinburgh, School of Mathematics, Edinburgh, United Kingdom.

used a log-Gaussian Cox process to model landslides and create spatial predictive maps to identify regions at risk in Italy. The data and the fitted intensity function were visualized using Google EarthTM; see Figure 1 and <https://vimeo.com/266344095>. The visualization of risk maps and how they evolve in time would highly benefit from immersive virtual reality (VR) tools, which more realistically render three-dimensional (3D) perspective.



Figure 1: Visualization of landslide data and fitted log-Gaussian Cox process intensity function using Google EarthTM; see also <https://vimeo.com/266344095>.

As explained by Castruccio *et al.*, and further motivated above, advanced 3D, immersive, or portable VR visualization techniques may be used to dynamically and interactively “*explore and assess the structure of the data and to improve resulting statistical models*”. We fully concur, although we want to point out that using such modern facilities as the KAUST 3D visualization cave, and creating smartphone or VR headset applications for scientific visualization often require a significant investment of time and the support from a specialized staff. Although this is undoubtedly worthwhile for the *communication of results* from statistical analyses, we think that for *data exploration*, simpler approaches such as R Shiny applications, or popular Geographic Information System (GIS) environments including Google EarthTM, which are visually efficient, cheaper, accessible, and significantly less time-demanding to set up, are still useful.

References

- Cressie, N. A. C. (1993) *Statistics for Spatial Data*. New York: Wiley.
- Davison, A. C. and Huser, R. (2015) Statistics of extremes. *Annual Review of Statistics and its Application* **2**, 203–235.
- Davison, A. C., Huser, R. and Thibaud, E. (2018) Spatial extremes. In *Handbook of Environmental and Ecological Statistics*, eds A. E. Gelfand, M. Fuentes and R. L. Smith. CRC Press. To appear.
- Davison, A. C., Padoan, S. and Ribatet, M. (2012) Statistical modelling of spatial extremes (with discussion). *Statistical Science* **27**(2), 161–186.
- Genton, M. G., Castruccio, S., Crippa, P., Dutta, S., Huser, R., Sun, Y. and Vettori, S. (2015) Visuanimation in statistics. *Stat* **4**(1), 81–96.
- Huser, R. and Davison, A. C. (2014) Space-time modelling of extreme events. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* **76**(2), 439–461.
- Lombardo, L., Opitz, T. and Huser, R. (2018) Point process-based modeling of multiple debris flow landslides using INLA: an application to the 2009 Messina disaster. *Stochastic Environmental Research and Risk Assessment* **32**, 2179–2198.